

Novocastra™ Liquid Mouse Monoclonal Antibody bcl-2 Oncoprotein

Product Code: NCL-L-bcl-2

Intended Use	FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES.
Specificity	Human bcl-2 oncoprotein.
Clone	bcl-2/100/D5
Ig Class	IgG1
Antigen Used for Immunizations	Synthetic peptide sequence (GAAPAGIFSSQPGC-COOH).
Hybridoma Partner	Mouse myeloma (p3-NS1-Ag4-1).
Preparation	Liquid tissue culture supernatant containing sodium azide. Volume as indicated on vial label.
Effective on Frozen Tissue	Yes
Effective on Paraffin Wax Embedded Tissue	Yes
Recommendations on Use	Immunohistochemistry on paraffin sections. Heat Induced Epitope Retrieval (HIER): Please follow the instructions for use in Novocastra Epitope Retrieval Solution pH 6. Suggested dilution: 1:100 for 30 minutes at 25 °C. This is provided as a guide and users should determine their own optimal working dilutions. Visualization: Please follow the instructions for use in the Novolink™ Polymer Detection Systems. For further product information or support, contact your local distributor or regional office of Leica Biosystems, or alternatively, visit the Leica Biosystems web site, www.LeicaBiosystems.com <u>The performance of this antibody should be validated when utilized with other manual staining systems or automated platforms.</u> Western Blotting: Typical working dilution 1:25–1:50.
Positive Controls	Immunohistochemistry: Tonsil. Western Blotting: Bristol 8 cell line.
Staining Pattern	Membrane and/or cytoplasmic.
Storage and Stability	Store at 2–8 °C. Do not freeze. Return to 2–8 °C immediately after use. Do not use after expiration date indicated on the vial label. Storage conditions other than those specified above must be verified by the user.
Warnings and Precautions	This reagent has been prepared from the supernatant of cell culture. As it is a biological product, reasonable care should be taken when handling it. This reagent contains sodium azide. A Material Safety Data Sheet is available upon request or available from www.LeicaBiosystems.com
General Overview	bcl-2 is an inner mitochondrial membrane protein that acts as an inhibitor of apoptosis. Apoptosis forms part of the normal pathway of B cell production. Consequently, bcl-2 expression is inhibited in the B cells of germinal centers unless they have been activated by signals which prevent their entry into apoptosis.





B I O S Y S T E M S

General References

Von Haefen C, Wieder T, Gillissen B et al. *Oncogene*. 2002; 21(25):4009-4019.
Takes RP, Baatenburg de Jong RJ, Wijffels K et al. *Journal of Pathology*. 2001; 194:298-302.
Tweddle DA, Malcolm AJ, Cole M et al. *American Journal of Pathology*. 2001; 158(6): 2067-2077.
Ramani P and Lu Q-L. *Journal of Pathology*. 1994; 172:273-278.
Pezzella F, Tse AGD, Cordell JL et al. *American Journal of Pathology*. 1990; 137(2):225-232.