

# Kreatech™ FISH probes

## Product Information Sheet

KBI-10305


RARA (17q21) Break



**Kreatech Biotechnology B.V.**  
Vierweg 20  
1032 LG Amsterdam  
The Netherlands  
[www.LeicaBiosystems.com](http://www.LeicaBiosystems.com)

PI-KBI-10305\_D1.1

Published March 2015



Not to scale

## Kreatech™ RARA (17q21) Break FISH probe

**Introduction:** Acute promyelocytic leukemia (APL) is mainly characterized by the PML-RARA t(15;17) translocation. But also other variants translocations involving RARA at 17q21 occur in APL, as t(11;17)(q23;q21), t(5;17)(q35;q12-21), t(11;17)(q13;q21), and t(17;17)(q21;q21), whereby RARA is fused to the PLZF, NPM1, NUMA1, and STAT5B genes, respectively,

**Intended use:** The **RARA (17q21) Break** FISH probe is optimized to detect translocations involving the RARA gene region at 17q21 in a dual-color, split assay on metaphase/interphase spreads.

The probe is recommended to be used in combination with one of the Kreatech Pretreatment kits providing necessary reagents to perform FISH on various sample types for optimal results. (see also [www.LeicaBiosystems.com](http://www.LeicaBiosystems.com) and look for Kits & reagents)

**Critical region 1 (red):** The **distal RARA** gene region probe is direct-labeled with PlatinumBright™550.  
**Critical region 2 (green):** The **proximal RARA** gene region probe is direct-labeled with PlatinumBright™495.

**Reagent:** Kreatech probes are direct-labeled DNA probes provided in a ready-to-use format. Apply 10 µl of probe to a sample area of approximately 22 x 22 mm.

**Please refer to the Instructions for Use for the entire Kreatech FISH protocol.**

**Kreatech FISH probes are REPEAT-FREE™ and therefore do not contain Cot-1 DNA. Hybridization efficiency is increased and background, due to unspecific binding, is highly reduced.**

**Interpretation:** The **RARA (17q21) Break** FISH probe is designed as a dual-color split probe to detect translocations at 17q21. A break is defined when a red/green or yellow fusion signal (F) splits into separate red and green signals. Only red and green signals which are more than one signal diameter apart from each other are counted as a break. Co-localized red/green or yellow signals identify the normal chromosome(s) 17. Signal patterns other than those described above may indicate variant translocations or other complex rearrangements. Investigators are advised to analyze metaphase cells for the interpretation of atypical signal patterns.

	Normal Signal Pattern	17q21 Split
Expected Signals	2F	1F1R1G

**References:** Grimwade et al, 2000, Blood 96; 1297-1308

**Warning and precautions:** In case of emergencies check SDS sheets for medical advice. SDS sheets may be obtained by either contacting Leica Technical Support or visiting [www.LeicaBiosystems.com](http://www.LeicaBiosystems.com). DNA probes contain formamide which is a teratogen; do not inhale or allow skin contact. Wear gloves and a lab coat when handling DNA probes. All materials should be disposed of according to your institution's guidelines for hospital waste disposal.

**Reagent Storage and Handling:** Store at 2-8 °C. Reagents should not be used after the expiration date on the vial label.

**TECHNICAL SUPPORT** Technical support is available at [www.LeicaBiosystems.com](http://www.LeicaBiosystems.com) or +31 20 6919181 or via e-mail: [kreatech-support@leicabiosystems.com](mailto:kreatech-support@leicabiosystems.com).

**CUSTOMER SERVICE** Kreatech probes may be ordered through Leica Customer Service +31 20 6919181 or order via e-mail: [purchase.orders@leica-microsystems.com](mailto:purchase.orders@leica-microsystems.com).