

Kreatech[™] FISH probes **Product Information Sheet**

KI-10737 FGFR1 (8p11) Break 100 µl

RUO - Research Use Only

Not for use in diagnostic procedures



RH62328 520 kB 8p11.2 RH7851 FGFR1 SHGC-103442 420 kB

SHGC-2804

Vlierweg 20

1032 LG Amsterdam The Netherlands

www.LeicaBiosystems.com

DANGER FORMAMIDE

8°C



Published Sept 2015

2°

Kreatech™ FGFR1 (8p11) Break FISH probe

Introduction: The FGFR1 (8p11) Break FISH probe is optimized to detect translocations involving the FGFR1 gene region at 8p11 in a dual-color.

Critical region 1 (red):The distal FGFR1 gene region probe is direct-labeled with PlatinumBright™550.Critical region 2 (green):The proximal FGFR1 gene region probe is direct-labeled with PlatinumBright™495.

Reagent:Kreatech probes are direct-labeled DNA probes provided in a ready-to-use format.
Apply 10 μl of probe to a sample area of approximately 22 x 22 mm.

Please refer to the Instructions for Use for the entire Kreatech FISH protocol.

Kreatech FISH probes are REPEAT-FREE[™] and therefore do not contain Cot-1 DNA. Hybridization efficiency is increased and background, due to unspecific binding, is highly reduced.

Patterns: The FGFR1 (8p11) Break FISH probe is designed as a dual-color split probe to detect translocations at 8p11. A break is defined when a red/green or yellow fusion signal (F) splits into separate red and green signals. Only (R) red and green (G) signals which are more than one signal diameter apart from each other are counted as a break. Co-localized red/green or yellow signals identify the normal chromosome(s) 8.

Signal patterns other than those described above may indicate variant translocations or other complex rearrangements. Investigators are advised to analyze metaphase cells for the interpretation of atypical signal patterns.

	Normal Signal Pattern	8p11 Split	8p11 Amplification
Expected Signals	2F	1F1R1G	3+F

References: Smedley et al, 1998, Hum Mol Genet. 7; 637-642.

Warning and precautions: In case of emergencies check SDS sheets for medical advice. SDS sheets may be obtained by either contacting Leica Technical Support or visiting <u>www.LeicaBiosystems.com</u>. DNA probes contain formamide which is a teratogen; do not inhale or allow skin contact. Wear gloves and a lab coat when handling DNA probes. All materials should be disposed of according to your institution's guidelines for hospital waste disposal.

Reagent Storage and Handling:	Store at 2-8 °C. Reagents should not be used after the expiration date on the vial label.
TECHNICAL SUPPORT	Technical support is available at www.LeicaBiosystems.com/service-support/technical-support/ or toll free at 800-248-0123 or via e-mail: keatech-support@leicabiosystems.com .
CUSTOMER SERVICE	Kreatech probes may be ordered through Leica Customer Service toll free at 800-248-0123 or order via e-mail: <u>purchase.orders@leica-microsystems.com</u> .