

Novocastra™ Liquid Mouse Monoclonal Antibody CD123

Product Code: NCL-L-CD123

Intended Use	FOR RESEARCH USE ONLY.
Specificity	Human CD123
Clone	BR4MS
Ig Class	IgG2b
Antigen Used for Immunizations	Prokaryotic recombinant protein corresponding to 101 amino acids of the external domain of the human CD123 molecule.
Hybridoma Partner	Mouse myeloma (p3-NS1-Ag4.1).
Preparation	Liquid tissue culture supernatant containing 15 mM sodium azide. Volume as indicated on vial label.
Effective on Frozen Tissue	Not evaluated.
Effective on Paraffin Wax Embedded Tissue	Yes (using heat induced epitope retrieval with Tris–EDTA-based buffer, pH 9.0: see overleaf)
Recommendations on Use	Immunohistochemistry: Typical working dilution 1:100. Heat induced epitope retrieval with Tris-EDTA–based buffer, pH 9.0. 30 minutes primary antibody incubation at 25 °C. Polymer detection recommended. Western Blotting: Not recommended.
Positive Controls	Immunohistochemistry: Tonsil
Staining Pattern	Cytoplasmic/membrane
Storage and Stability	Store liquid antibody at 4 °C. Under these conditions, there is no significant loss in product performance up to the expiry date indicated on the vial label. Prepare working dilutions on the day of use.
General Overview	The CD123 antigen is also known as the alpha subunit of the human interleukin-3 receptor. It is a type I transmembrane glycoprotein and is a member of the cytokine receptor superfamily. CD123 forms a heterodimer with CD131 (the beta subunit of the interleukin-3 receptor) to form the interleukin-3 receptor, where the cytokine specificity is provided by the alpha subunit and the signal transduction function is provided by the beta subunit. The interleukin-3 receptor is reported to be expressed on monocytes, neutrophils, basophils, eosinophils, megakaryocytes, erythroid precursors, mast cells, macrophages and a subpopulation of B cells, where it mediates proliferation and differentiation of these cells. Outside the hematopoietic system CD123 is reported to be expressed in Leydig cells of the testis, some endothelial cells, and cells of the placenta and brain.
General References	Garnache-Ottou F, Feuillard J and Saas P. British Journal of Haematology. 2007; 136:539–548. Moretti S, Lanza F and Dabusti M et al. Journal of Biological Regulators and Homeostatic Agents. 2001; 15:98–100.



Instructions for Use

Heat Induced Epitope Retrieval Combined With Polymer Detection For Immunohistochemical Demonstration On Paraffin Sections

1. Cut and mount sections on slides coated with a suitable tissue adhesive.
2. Deparaffinize sections and rehydrate to distilled water.
3. Place sections in 0.5% hydrogen peroxide/methanol for 10 minutes (or use other appropriate endogenous peroxidase blocking procedure). Wash sections in tap water.
4. Heat 1500 mL of the recommended epitope retrieval solution (Citrate based pH 6.0 - Epitope Retrieval Solution unless otherwise indicated overleaf) in a stainless steel pressure cooker until boiling. Cover but do not lock lid.
5. Position slides into metal staining racks (do not place slides close together as uneven staining may occur) and lower into pressure cooker ensuring slides are completely immersed in epitope retrieval solution. Lock lid.
6. When the pressure cooker reaches operating temperature and pressure (after about 5 minutes) start a timer for 1 minute (unless otherwise indicated on the data sheet).
7. When the timer rings, remove pressure cooker from heat source and run under cold water with lid on. **DO NOT OPEN LID UNTIL THE INDICATORS SHOW THAT PRESSURE HAS BEEN RELEASED.** Open lid, remove slides and place immediately into a bath of tap water.
8. Wash sections once using fresh Tris-Buffered Saline (TBS, pH 7.6) buffer for 5 minutes.
9. Place sections in diluted normal serum (eg NCL-G-SERUM) for 10 minutes.
10. Incubate sections with primary antibody.
11. Wash twice, each time using fresh TBS buffer for 5 minutes.
12. For visualization of the bound primary antibody, follow instructions supplied with the Polymer Detection System.
13. Counterstain with hematoxylin (if required), dehydrate and mount.

* (In most applications, Phosphate Buffered Saline, pH 7.6, can be used instead of TBS, pH 7.6).

Safety Note

To ensure the correct and safe use of your pressure cooker, PLEASE READ THE MANUFACTURER'S INSTRUCTIONS.