

Novocastra™ Liquid Mouse Monoclonal Antibody Cytokeratin 7

Product Code: NCL-L-CK7-OVTL

Intended Use	FOR RESEARCH USE ONLY.
Specificity	Human cytokeratin 7 intermediate filament protein.
Clone	OV-TL 12/30
Ig Class	IgG1, kappa
Antigen Used for Immunizations	OTN 11 ovarian carcinoma cell line.
Hybridoma Partner	Mouse myeloma (Sp2/0-Ag14).
Preparation	Liquid tissue culture supernatant containing 15 mM sodium azide. Volume as indicated on vial label.
Effective on Frozen Tissue	Yes. Acetone fixation recommended.
Effective on Paraffin Wax Embedded Tissue	Yes (using the high temperature antigen unmasking technique: see overleaf, or trypsin digestion of sections).
Recommendations on Use	Immunohistochemistry: Typical working dilution 1:50. High temperature antigen unmasking technique, or trypsin digestion of sections. 60 minutes primary antibody incubation at 25 °C. Standard ABC technique. Western Blotting: Typical working dilution 1:500–1:1000.
Positive Controls	Immunohistochemistry: Breast. Western Blotting: HeLa cell line.
Staining Pattern	Cytoplasmic and membrane.
Storage and Stability	Store liquid antibody at 4 °C. Under these conditions, there is no significant loss in product performance up to the expiry date indicated on the vial label. Prepare working dilutions on the day of use.
General Overview	Clone OV-TL 12/30 reacts with the human cytokeratin intermediate filament protein (54 kD) identified as cytokeratin 7. It reacts with a large number of epithelial cell types including many ductal and glandular epithelia and distinguishes between certain types of normal glandular epithelia of which lung and breast are positive and colon and prostate are negative. No reactivity with other cytokeratins has been observed and, in general, NCL-L-CK7-OVTL does not react with stratified squamous epithelia but it does react with transitional epithelium of the urinary tract. Hepatocytes are negative while bile duct epithelial cells are positive.
General References	Torenbek R, Lagendijk J H, Van Diest P J, et al.. <i>Histopathology</i> . 32: 20–27 (1998). van de Molengraft F J J M, van Niekerk C C, Jap P H K, et al.. <i>Histopathology</i> . 22: 35–38 (1993). Markey A C, Lane E B, MacDonald D M, et al.. <i>British Journal of Dermatology</i> . 126: 154–160 (1992). Ramaekers F, Huysmans A, Schaart G, et al.. <i>Experimental Cell Research</i> . 170: 235–249 (1987). Osborn M, van Lessen G, Weber K, et al.. <i>Laboratory Investigation</i> . 55 (4): 497–504 (1986). Moll R, Franke W W, Schiller D L. <i>Cell</i> . 31: 11–24 (1982).



Instructions for Use



High Temperature Antigen Unmasking Technique Followed by Trypsin Digestion for Immunohistochemical Demonstration on Paraffin Sections

1. Cut and mount sections on slides coated with a suitable tissue adhesive.
2. Deparaffinize sections and rehydrate to distilled water.
3. Place sections in 0.5% hydrogen peroxide/methanol for 10 minutes (or use other appropriate endogenous peroxidase blocking procedure). Wash sections in tap water.
4. Heat 1500 mL of the recommended unmasking solution (0.01 M citrate buffer, pH 6.0 (or Epitope Retrieval Solution, RE7113) unless otherwise indicated overleaf - see other Epitope Retrieval Solutions in the range) until boiling in a stainless steel pressure cooker. Cover but do not lock lid.
5. Position slides into metal staining racks (do not place slides close together as uneven staining may occur) and lower into pressure cooker ensuring slides are completely immersed in unmasking solution. Lock lid.
6. When the pressure cooker reaches operating temperature and pressure (after about 5 minutes) start a timer for 1 minute (unless otherwise indicated on the data sheet).
7. When the timer rings, remove pressure cooker from heat source and run under cold water with lid on. DO NOT OPEN LID UNTIL THE INDICATORS SHOW THAT PRESSURE HAS BEEN RELEASED. Open lid, remove slides and place immediately into a bath of tap water.
8. Place slides in pre-heated distilled water to bring the sections to 37 °C for a minimum of 5 minutes.
9. Incubate sections in pre-heated Trypsin solution at 37 °C for 30 seconds.
10. Rinse sections in running tap water.
11. Proceed with immunohistochemistry protocol.
12. Wash sections in TBS* buffer (pH 7.6) for 1 x 5 minutes.
13. Place sections in diluted normal serum (or RTU Normal Horse Serum) for 10 minutes.
14. Incubate sections with primary antibody. Use Antibody Diluent RE7133 (where available).
15. Wash in TBS buffer for 2 x 5 minutes.
16. Incubate sections in an appropriate biotinylated secondary antibody.
17. Wash in TBS buffer for 2 x 5 minutes.
18. Incubate slides in ABC reagent (or RTU streptavidin/peroxidase complex).
19. Wash in TBS buffer for 2 x 5 minutes.
20. Incubate slides in DAB or other suitable peroxidase substrate.
21. Wash thoroughly in running tap water.
22. Counterstain with hematoxylin (if required), dehydrate and mount.

Solutions

Trypsin Solution

**Trypsin containing chymotrypsin should always be used. The enzyme activities can vary from a supplier and between batches. Such variations may affect the incubation time required.*

Preheat the following to 37 °C using a water bath:

- (i) 200 mL of TBS
- (ii) 200 mL of distilled water.

Dissolve 0.2 g Trypsin 250 and 0.2 g Calcium Chloride in the 200 mL of TBS.

Once the Trypsin solution is at 37 °C, pH to 7.8 with 1 M sodium hydroxide.

0.01 M Citrate Buffer (pH 6.0) or RE7113 (where available).

Add 3.84 grams of Citric acid (anhydrous) to 1.8 L of distilled water. Adjust to pH 6.0 using concentrated NaOH. Make up to 2 L with distilled water.

1 mM EDTA (pH 8.0) or RE7116 (where available)

Add 0.37 g of EDTA (SIGMA product code E-5134) to 1 L of distilled water. Adjust pH to 8.0 using 1.0 M NaOH.

20 mM TRIS/0.65 mM EDTA/0.0005% TWEEN (pH 9.0) or RE7119 (where available)

Dissolve 14.4 g Tris (BDH product code 271197K) and 1.44 g EDTA (SIGMA product code E-5134) to 0.55 L of distilled water. Adjust pH to 9.0 with 1 M HCl and add 0.3 mL Tween 20 (SIGMA product code P-1379). Make up to 0.6 L with distilled water. This is a 10x concentrate which should be diluted with distilled water as required (eg 150 mL diluted with 1350 mL of distilled water).

** In most applications, 10mM phosphate, 0.15 M NaCl, pH 7.6 (PBS) can be used instead of 50 mM Tris, 0.15 M NaCl, pH 7.6 (TBS).*

Safety Note

To ensure the correct and safe use of your pressure cooker, PLEASE READ THE MANUFACTURER'S INSTRUCTIONS.

protocol/HTAUT + TRYP SIN/07/08